

## Review

# Drought stress and reactive oxygen species

## Production, scavenging and signaling

Maria Helena Cruz de Carvalho

Laboratoire d'Ecophysiologie Moléculaire; UMR IRD 137 BioSol; Université Paris 12; Créteil, France

**Abbreviations:** ABA, abscisic acid; AOX, alternative oxidase; APX, ascorbate peroxidase; ASC, ascorbate; CAT, catalase; EST, expressed sequence tags; GPX, glutathione peroxidase; GR, glutathione reductase; GSH, reduced glutathione; PEG, polyethyleneglycol; PSI, PSII, photosystems I and II; ROS, reactive oxygen species; RuBP, ribulose-1,5-bisphosphate; rubisco, ribulose-1,5-bisphosphate carboxylase/oxidase; SOD, superoxide dismutase

**Key words:** drought stress, stomatal closure, oxidative stress, Mehler reaction, water-water cycle, photorespiration, scavenging enzymes, ROS signaling, H<sub>2</sub>O<sub>2</sub>, ABA

As sessile organisms, plants have evolved mechanisms that allow them to adapt and survive periods of drought stress. One of the inevitable consequences of drought stress is enhanced ROS production in the different cellular compartments, namely in the chloroplasts, the peroxisomes and the mitochondria. This enhanced ROS production is however kept under tight control by a versatile and cooperative antioxidant system that modulates intracellular ROS concentration and sets the redox-status of the cell. Furthermore, ROS enhancement under stress functions as an alarm signal that triggers acclimatory/defense responses by specific signal transduction pathways that involve H<sub>2</sub>O<sub>2</sub> as secondary messenger. ROS signaling is linked to ABA, Ca<sup>2+</sup> fluxes and sugar sensing and is likely to be involved both upstream and downstream of the ABA-dependent signaling pathways under drought stress. Nevertheless, if drought stress is prolonged over to a certain extent, ROS production will overwhelm the scavenging action of the antioxidant system resulting in extensive cellular damage and death.

### Introduction

Drought is one of the most important manifestations of abiotic stress in plants. It is the major yield-limiting factor of crop plants and it actively and continuously determines the natural distribution of plant species. Drought exacerbates the effect of the other stresses to which plants are submitted (abiotic or biotic) and several different abiotic stresses result in water stress (like salt and cold stresses). As sessile organisms, plants have to cope with drought stress at least at some point in their life cycle. They have however evolved mechanisms that allow them to adapt and survive periods of water deficit, if not at the whole plant level, at some level or form of plant structure.

According to the type of strategy adopted, plants are said to escape, avoid or tolerate drought stress,<sup>1</sup> although these are not mutually exclusive. The plant drought response will depend on the species inherent "strategy" but also on the duration and severity of the drought period. If prolonged over to a certain extent drought stress will inevitably result in oxidative damage due to the over production of reactive oxygen species.<sup>2</sup>

Reactive oxygen species (ROS), also called active oxygen species (AOS) or reactive oxygen intermediates (ROI) are the result of the partial reduction of atmospheric O<sub>2</sub>. There are basically four forms of cellular ROS, singlet oxygen (<sup>1</sup>O<sub>2</sub>), superoxide radical (O<sub>2</sub><sup>-</sup>), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and the hydroxyl radical (HO<sup>•</sup>), each with a characteristic half-life and an oxidizing potential. ROS can be extremely reactive, especially singlet oxygen and the hydroxyl radical and, unlike atmospheric oxygen, they can oxidize multiple cellular components like proteins and lipids, DNA and RNA. Unrestricted oxidation of the cellular components will ultimately cause cell death.<sup>3</sup>

Unlike biotic stress where an oxidative burst is part of a defense response that frequently triggers programmed cell death (PCD),<sup>4</sup> the role of ROS production and control during drought stress is yet to be resolved. However, as stated by Dat and collaborators,<sup>5</sup> ROS seem to have a dual effect under abiotic stress conditions that depend on their overall cellular amount. If kept at relatively low levels they are likely to function as components of a stress-signaling pathway, triggering stress defense/acclimation responses.<sup>5,6</sup> However, when reaching a certain level of phytotoxicity ROS become extremely deleterious, initiating uncontrolled oxidative cascades that damage cellular membranes and other cellular components resulting in oxidative stress and eventually cell death.<sup>3,5</sup> This review will focus on ROS production and control during drought stress with highlights on the signaling aspects of ROS generation and action.

### Plants are Natural Producers of ROS

Under normal conditions plants are continuously producing ROS. Unlike singlet oxygen and the hydroxyl radical whose production is kept at minimum levels,<sup>7,8</sup> superoxide and H<sub>2</sub>O<sub>2</sub> are synthesized at

Correspondence to: Maria Helena Cruz de Carvalho; Laboratoire d'Ecophysiologie Moléculaire; UMR IRD 137 BioSol; Université Paris 12; 61 avenue du Général de Gaulle; F- 94010 Créteil Cedex France; Tel.: + 33.1.45.17.65.65; Fax: +33.1.45.17.16.17; E-mail address: mhcc@univ-paris12.fr

Submitted: 01/08/08; Accepted: 02/26/08

Previously published online as a *Plant Signaling & Behavior* E-publication: [www.landesbioscience.com/journals/psb/article/5537](http://www.landesbioscience.com/journals/psb/article/5537)

very high rates under normal conditions.<sup>9</sup> One of the major cellular sites responsible for ROS production is the chloroplast.<sup>10</sup> During photosynthesis, energy from the sunlight is captured and transferred to two light-harvesting complexes (photosystem II and photosystem I) in the chloroplast thylakoidal membranes. A succession of redox reactions occurs within the electron transport chain in the light, until electrons finally reach  $\text{CO}_2$ , in the dark reactions. However, it is not uncommon that through this path other final acceptors of electrons are used, namely oxygen.<sup>11</sup> Singlet oxygen can be formed by energy transfer from triplet excited state chlorophyll to  $\text{O}_2$ .<sup>12</sup> On the other hand, the thylakoidal electron transport components on the PSI side such as the Fe-S centers and the reduced thiredoxin are auto-oxidizable resulting in the reduction of  $\text{O}_2$  (the Mehler reaction) thus forming superoxide and  $\text{H}_2\text{O}_2$ .<sup>12,13</sup> It has been estimated that approximately 10% of the photosynthetic electrons flow to the Mehler reaction.<sup>14</sup> This “leakage” of electrons to  $\text{O}_2$  with the generation of ROS is in fact favorable to the electron transport chain since it poises electron carriers thus making them more efficient.<sup>9</sup>

During photosynthesis there is a different pathway, called photorespiration that can also generate ROS (Fig. 1). In fact, rubisco, the enzyme that catalyses the carboxylation of ribulose-1,5-bisphosphate (RuBP) during carbon assimilation, can also use  $\text{O}_2$  to oxygenate ribulose-1,5-bisphosphate. This reaction yields glycolate that is then transported from chloroplasts to peroxisomes where they are oxidized by glycolate oxidase and  $\text{H}_2\text{O}_2$  is generated.<sup>15</sup>

Mitochondrial electron transport chain is also responsible for ROS generation under normal conditions, although to a lesser extent than chloroplasts and peroxisomes in the light.<sup>16</sup> It has been estimated that approximately 1–2% of the consumed oxygen, respired by plant mitochondria, will be used to form superoxide.<sup>17</sup> This ROS production is likely to occur mainly in complexes I and III of the mitochondrial electron transport chain.<sup>18</sup>

## Plants Keep ROS Under Control by an Efficient and Versatile Scavenging System

In order to cope with continuous ROS production plants have a battery of enzymatic and nonenzymatic antioxidants, which function as an extremely efficient cooperative system. The major scavenging mechanisms include superoxide dismutase (SOD), enzymes and metabolites from the ascorbate-glutathione cycle, and catalase (CAT).<sup>9,19,20</sup> They are located throughout the different compartments of the plant cell, with the exception of catalase that is exclusively located in peroxisomes. SOD is the front-line enzyme in ROS attack since it rapidly scavenges superoxide, one of the first ROS to be produced, dismutating it to oxygen and  $\text{H}_2\text{O}_2$ .<sup>19</sup> However, this reaction only converts one ROS to another, and  $\text{H}_2\text{O}_2$  also needs to be destroyed since it promptly attacks thiol proteins.<sup>9</sup> The major enzymatic cellular scavengers of  $\text{H}_2\text{O}_2$  are catalase and ascorbate peroxidase (APX).<sup>9,20</sup> They have however different affinities for this ROS and seem to have different cellular roles in  $\text{H}_2\text{O}_2$  scavenging. In fact CAT does not need a reductant to scavenge  $\text{H}_2\text{O}_2$  making it reducing power-free, whereas APX needs a reductant, ascorbate. On the other hand, CAT has a lower affinity for  $\text{H}_2\text{O}_2$  (mM range) than APX ( $\mu\text{M}$  range).<sup>3</sup> All this gathered has led to the hypothesis that APX, an enzyme located in every cellular ROS producing compartment, might function as a fine regulator of intracellular ROS steady-state levels, possibly for signaling purposes, whereas CAT,




Figure 1. The photorespiratory pathway in photosynthetic plants cells. This cycle involves the oxygenation of RuBP in the chloroplast (1), the formation of  $\text{H}_2\text{O}_2$  in the peroxisomes (2) and the conversion of glycine to serine in the mitochondria (3). RuBP, ribulose-1,5-bisphosphate.

located exclusively in the peroxisomes, might function as a bulk remover of excess ROS production under stress conditions.<sup>3,9,20</sup>

Glutathione reductase (GR), the last enzyme of the ascorbate/glutathione cycle, has a major role in maintaining the intracellular glutathione pool in the reduced state (GSH).<sup>9</sup> GSH can function as an antioxidant either directly (non enzymatically), like ascorbate, by scavenging singlet oxygen, superoxide or even hydroxyl radicals, or indirectly as a reducing agent that recycles ascorbic acid from its oxidized form to its reduced form by the enzyme dehydroascorbate reductase.<sup>9,21</sup>

In the chloroplast, the Mehler reaction occurring during photosynthesis is an important alternative sink for electrons, but it produces superoxide as side effect. This active oxygen species is however rapidly dismutated by a membrane bound superoxide dismutase (SOD), producing  $\text{H}_2\text{O}_2$ .  $\text{H}_2\text{O}_2$  is then locally converted to water by ascorbate peroxidase (APX). This is called the Mehler-peroxidase reaction or the water-water cycle.<sup>22</sup> One of the major advantages of the water-water cycle is the scavenging of superoxide and  $\text{H}_2\text{O}_2$  at their production site, without further damage to the thylakoids or other cellular compartments (Fig. 2). Furthermore, the water-water cycle seems to be autonomous regarding its energy supply since it gets its reducing power directly from the photosynthetic apparatus.<sup>22</sup>

$\text{H}_2\text{O}_2$  is a very stable ROS with the longest half-life (~1 ms) and is also the more diffusive<sup>4</sup> so it can readily “escape” from the organelle where it was produced to the cytosol. It can also be directly produced in the cytosol by existing cytosolic SODs. However in the cytosol,  $\text{H}_2\text{O}_2$  is readily scavenged by ascorbate through the ascorbate/glutathione cycle.

## ROS Production is Enhanced Under Drought Stress

The first plant organ to detect a limitation on the water supply is the root system. It has been shown that besides water and minerals, roots also send signals to the leaves through the xylem sap, and the phytohormone abscisic acid is considered to be one of the major root-to-shoot stress signals.<sup>23–26</sup> When the stress signal reaches




Figure 2. The water-water cycle or the Mehler-peroxidase reaction involves the leakage of electrons from the photosynthetic electron transport chain to oxygen with the generation of superoxide (1), which is further dismutated by SOD forming  $\text{H}_2\text{O}_2$  (2).  $\text{H}_2\text{O}_2$  is then scavenged by a thylakoidal APX with the generation of water (3). This cycle gets its reducing power through ascorbate regeneration by ferredoxin (4). Side reactions that can occur in the chloroplasts are the metal catalyzed Haber-weiss/Fenton reactions that result in the formation of the highly reactive hydroxyl radical (5). PSI, photosystem I, Fd, ferredoxin, MDA, monodehydroascorbate, AA, ascorbate, tAPX, thylakoidal ascorbate peroxidase, SOD, superoxide dismutase.

the leaves, it triggers stomatal closure and the plant shifts to a water-saving strategy. Hence, by adjusting stomatal opening, plants are able to control water loss by reducing the transpiration flux, but they are concomitantly limiting the entrance of carbon dioxide ( $\text{CO}_2$ ). This will have direct and indirect effects on the reduction of net photosynthesis and on the overall production of ROS by plants under drought stress.<sup>3</sup> There are many studies that report on increased ROS accumulation and oxidative stress under drought stress.<sup>27-32</sup> In fact, under drought stress, ROS production is enhanced through multiple ways. For instance, the limitation on  $\text{CO}_2$  fixation will reduce  $\text{NADP}^+$  regeneration through the Calvin cycle, hence provoking an over reduction of the photosynthetic electron transport chain. In fact, during photosynthesis and under drought stress there is a higher leakage of electrons to  $\text{O}_2$  by the Mehler reaction.<sup>2</sup> It was estimated that in drought stressed wheat, the leakage of photosynthetic electrons to the Mehler reaction was increased by approximately 50% as compared to unstressed wheat.<sup>33</sup> An increase on thylakoid membrane electron leakage to  $\text{O}_2$  under drought stress was also seen in sunflower.<sup>34</sup> However, it is quite difficult to assess the part of ROS generated by the Mehler reaction to that generated by photorespiration. In fact, under drought stress the photorespiratory pathway is also enhanced, especially when RuBP oxygenation is maximal due to limitation on  $\text{CO}_2$  fixation.<sup>35</sup> The predominance of photorespiration on the oxidative load under drought stress has been recently put forward. Noctor and collaborators<sup>35</sup> have estimated that photorespiration is likely to account for over 70% of total  $\text{H}_2\text{O}_2$  production under drought stress conditions.

The chloroplast is a quite robust cellular compartment towards ROS because of the different scavenging enzymes and metabolites present.<sup>22,35,36</sup> However, under drought stress one of the real threats towards the chloroplast is the production of the hydroxyl radical in the thylakoids through “iron-catalysed” reduction of  $\text{H}_2\text{O}_2$  by both SOD and ascorbate. The hydroxyl radical is the ROS which has the shortest half-life ( $\sim 1 \mu\text{s}$ ) but it also has an extremely strong oxidizing potential reacting with almost every biological molecule.<sup>5</sup> Furthermore, there is no enzymatic reaction known to eliminate the highly reactive hydroxyl radical<sup>6,37</sup> and its accumulation will inevitably lead to deleterious reactions which damage the thylakoidal membranes and the photosynthetic apparatus.

## ROS Scavenging and Protection Under Drought

Plants can use the level of steady-state cellular ROS to monitor their intracellular level of stress.<sup>3</sup> However, this steady-state level must be tightly regulated in order to prevent an oxidative burst by over accumulation of ROS, which would ultimately result in extensive cell damage and death.<sup>3,5,12,22</sup> Symptoms of oxidative damage (like lipid peroxidation) have been used to assess the increase in ROS production under drought stress. However, the lack of symptoms does not imply that increased ROS formation is not occurring.<sup>2</sup> Instead, the lack of symptoms is likely to result on the concomitant increase in cellular antioxidant defenses.

**Scavenging enzymes.** The measure of specific antioxidant enzyme activities and/or expression analysis during water stress treatments has been generally accepted as an approach to assess the involvement of the scavenging system during drought stress. However, contradictory results have been gathered through the years (Table 1). These differences might be related to the plant age and tolerance/strategy towards water stress, but also to the duration and the intensity of the stress treatment. Nevertheless, some authors have detected a direct correlation between the level of the induction of the antioxidant system and the degree of drought tolerance of the plant species (same genus) or the plant cultivar (same species).<sup>31,38-47</sup>

In sunflower seedlings and in grass plants (*Aegilops squarrosa*) a decrease in SOD activity was detected under water stress.<sup>48,49</sup> Opposite to these results were those obtained in wheat,<sup>49</sup> in pea,<sup>50</sup> in common and tepary bean,<sup>46</sup> rice<sup>51</sup> and in olive trees,<sup>52</sup> where water stress increased SOD activity. Lower SOD activity might translate a lower production of superoxide by the Mehler reaction by keeping slight stomatal opening, thus avoiding complete inhibition of  $\text{CO}_2$  fixation.<sup>46</sup> Early regulation of stomatal conductance without complete stomatal closure is a characteristic strategy of drought adapted plants such as cowpea<sup>53</sup> and tepary bean.<sup>46</sup> Over expression of SOD in transgenic plants has given quite satisfactory results towards increasing oxidative stress tolerance.<sup>54-59</sup> It could be suggested that enhancement of oxidative stress tolerance is likely to have a positive effect on drought stress tolerance. Confirming this, two independent works have shown that transgenic alfalfa<sup>60</sup> and transgenic rice plants<sup>61</sup> over-expressing a chloroplast targeted MnSOD were more drought tolerant than the wild type plants. Furthermore, the transgenic rice plants over-expressing a pea MnSOD under the control of an oxidative stress inducible promoter, presented an enhanced photosynthetic capacity under a PEG imposed drought treatment.<sup>61</sup> This led the authors to suggest a role for chloroplastic SOD in ROS scavenging during drought stress, eventually through the water-water cycle, hence protecting the photosynthetic apparatus from drought-induced oxidative damage.<sup>61</sup>

Regarding the two major enzymes of the ascorbate/glutathione scavenging pathway, it was shown that APX and/or GR activities were enhanced during drought stress in cotton and spurred anoda,<sup>62</sup> wheat seedlings,<sup>63</sup> beans,<sup>46,47</sup> the moss *Tortula ruralis*,<sup>64</sup> rice,<sup>51</sup> alfalfa<sup>65</sup> and in cowpea.<sup>39,40</sup> A time course measure of APX and GR activities under a mild water stress imposed by a PEG treatment ( $-0,7 \text{ MPa}$ ) on maize detached leaves also showed a significant increase in both activities.<sup>66</sup> However, in pea plants the enzymes of the ascorbate/glutathione cycle showed a slight decreased activity under moderate drought stress (leaf  $\Psi_w = -1,30 \text{ MPa}$ ) followed by a 50% decreased activity under severe drought stress (leaf  $\Psi_w =$

Table 1 Drought stress responses of the major plant scavenging enzymes\*

Drought responses	Plant source	Refs	
<b>Scavenging enzyme activity increase</b>			
SOD	Wheat	49,139	
	Wheat <sup>T</sup>	42	
	Pea	50	
	Common bean	46	
	Tepary bean	46	
	Olive tree	52	
	Rice	51	
	Rice <sup>T</sup>	41	
	<i>Populus przewalskii</i>	140	
	<i>Coffea canephora</i>	141	
	<i>Festuca arundinacea</i> Schreb.	75	
	<i>Poa pratensis</i> L.	75	
	GR	Cotton	62
		Spurred anoda	62
Wheat <sup>S</sup>		31	
Wheat <sup>T</sup>		42	
Cowpea <sup>S</sup>		39	
Common bean		46,47	
Tepary bean		46	
<i>Tortula ruralis</i>		64	
Maize		66	
<i>Populus przewalskii</i>		140	
<i>Coffea canephora</i>		45,141	
Spring wheat		142	
Poplar <sup>T</sup>		44	
APX		Rice	51
	Rice <sup>T</sup>	41	
	Cotton	62	
	Spurred anoda	62	
	Cowpea <sup>S</sup>	40	
	Maize	66	
	Pea	67	
	Tepary bean	46	
	<i>Populus przewalskii</i>	140	
	<i>Coffea canephora</i>	45,141	
	Wheat <sup>T</sup>	42	
	Poplar <sup>T</sup>	44	
	Alfalfa	65	
	GPX/GST	<i>Tortula ruralis</i>	64
Rice <sup>T</sup>		41	
CAT	Alfalfa	65	
	Pea	50	
	Maize	66	
	Wheat	73	
	<i>Coffea canephora</i>	45,141	
Rice <sup>T</sup>	41		
<b>Scavenging enzyme activity decrease/unchanged</b>			
SOD	Wheat <sup>S</sup>	42	
	Sunflower	48	
	<i>Aegilops squarrosa</i>	49	
	Rice	41	
	Alfalfa	65	
GR	Wheat <sup>S</sup>	42	
	Pea	67	
	Cowpea <sup>T</sup>	39	
	Common bean <sup>S</sup>	47	
APX	Wheat <sup>S</sup>	42	
	Pea	67	
	Cowpea <sup>T</sup>	40	
	Rice	41	
GPX/GST	Spring wheat	142	
	Wheat <sup>S</sup>	42	
CAT	Tepary bean	46	
	Common bean	46	
	Sunflower	74	
	Sorghum	74	
	Rice	41,51	
	<i>Festuca arundinacea</i> Schreb.	75	
	<i>Poa pratensis</i> L.	75	

\* Abbreviations: SOD, superoxide dismutase; GR, glutathione reductase; APX, ascorbate peroxidase; GPX, glutathione peroxidase; GST, glutathione S-transferase; CAT, catalase; T, drought tolerant cultivar; S, drought susceptible cultivar.

-1,93 MPa).<sup>67</sup> The authors argued that this was probably due to the reduction on NADPH regeneration by photosynthesis under severe drought stress (which was shown to decrease by 70%), as well as to substrate depletion (GSH and ASC). Increased transcript accumulation of cytosolic APX was detected under drought stress in pea plants and this was accompanied by a slight enhanced APX protein content and APX activity.<sup>50</sup> Due to the high affinity of APX towards H<sub>2</sub>O<sub>2</sub>, this “small” increase on APX activity under drought stress was nonetheless suggested to be responsible for the scavenging of the elevated intracellular levels of H<sub>2</sub>O<sub>2</sub> produced under this stress condition.<sup>50</sup> The measure of total anti-oxidative enzyme activities does not account for what occurs in the different cellular compartments under drought stress and much information is being missed. In cowpea leaves it was shown that subtle changes occurred in the intracellular distribution of the APX and GR isoenzymes in response to a progressive drought stress that could relate to the tolerance of the cultivar.<sup>39,40</sup>

Recently, in an attempt to raise abiotic stress tolerance, simultaneous over-expression of both APX and CuZnSOD enzymes in transgenic potato and in tall fescue plants has been shown to result in an increased chloroplastic ROS scavenging action.<sup>68,69</sup> These works show that the combination of APX and CuZnSOD expression under the control of an oxidative stress-induced promoter, with simultaneous targeting to chloroplasts, proves to be quite efficient in increasing tolerance to abiotic stress-induced oxidative damage.<sup>68,69</sup>

Besides GR, two other enzymes related to glutathione metabolism, glutathione S-transferase, (GST) and glutathione peroxidase (GPX), have been shown to be induced under drought stress in the moss *Tortula ruralis*<sup>64</sup> and in leafy spurge (*Euphorbia esula*).<sup>70</sup> Glutathione peroxidases (GPXs) are efficient scavengers of H<sub>2</sub>O<sub>2</sub> and lipid hydroperoxides using GSH as a reducing agent but it has been suggested that in plants they preferably use thioredoxin as a reductant.<sup>71,72</sup> Hence, in a tight relationship with thioredoxins, plant GPXs are likely to be effective protectants of biomembranes under oxidative stress conditions.

Reports on catalase activity under drought stress are also heterogeneous. CAT activity has been shown to increase<sup>50,65,66,73</sup> and also to remain unchanged or even decrease under water stress.<sup>46,74,75</sup> Luna et al<sup>73</sup> have shown that severe drought stress induced an enhancement of CAT activity but this was not related to enhanced transcript accumulation, hence suggesting a rather more complex system of regulation. Transgenic tobacco plants expressing a bacterial catalase were shown to be photosynthetically more tolerant to high irradiance during drought stress than the wild type.<sup>76</sup> Furthermore, these authors have suggested that CAT is a less susceptible scavenging enzyme than APX regarding oxidative stress. Taken together these different reports seem to indicate that CAT activity is only enhanced under severe drought stress whereas under moderate drought stress H<sub>2</sub>O<sub>2</sub> scavenging is preferably made by ascorbic acid through the ascorbate/glutathione cycle. CAT has in fact a lower affinity for H<sub>2</sub>O<sub>2</sub> than APX which suggest its role in counteracting excessive H<sub>2</sub>O<sub>2</sub> production. Furthermore, excess H<sub>2</sub>O<sub>2</sub> may attack and inhibit APX, hence CAT activity is likely to be favorable in maintaining APX activity under severe drought stress.

Interesting results have shown that double antisense tobacco plants lacking both APX and CAT activate an alternative/redundant defense mechanism that compensates for the lack of these antioxidant




Figure 3. Proposed model of the drought stress response in three successive phases. The normal ROS steady-state level is disturbed by drought stress (I). Enhancement on ROS production due to stomatal closure shifts the equilibrium upwards and this triggers defense signal transduction pathways (II), prolonged drought stress will result in exacerbated ROS production that cannot be counterbalanced by the antioxidant system, leading to deleterious oxidative events which ultimately result in cell death (III).

enzymes.<sup>77</sup> Amongst the mechanisms that correlated to reduced susceptibility to oxidative stress of the double antisense plants was suppressed photosynthetic activity, the induction of metabolic genes of the pentose phosphate pathway, the induction of monodehydroascorbate reductase, and the induction of a chloroplastic homologue of the mitochondrial alternative oxidase (*IMMUTANS* gene).<sup>77</sup> These authors showed that a similar mechanism was not activated in single antisense plants lacking APX or CAT, which intriguingly rendered these plants more sensitive to oxidative stress compared to double antisense plants. This highlights the plasticity of the plant genome which is able to adjust and compensate for the lack of one or more scavenging enzymes under oxidative stress conditions.<sup>77</sup>

**Other antioxidants.** Superoxide produced by the Mehler reaction can also be directly reduced by ascorbate which is present at high concentrations in the chloroplast.<sup>9</sup> Apart from ascorbate and glutathione, plant cells possess other non enzymatic antioxidants that also participate in ROS scavenging or avoidance under normal and drought stress conditions. Their accumulation under drought stress relates to the drought tolerance of the plant species.<sup>78</sup> Alpha tocopherol is an antioxidant that not only prevents the formation of singlet oxygen and the hydroxyl radicals, but also scavenges lipid peroxyl radicals.<sup>78</sup> Under drought stress, this potent protector of thylakoidal and chloroplastic membranes has been shown to accumulate in several plant species.<sup>63,78,79</sup> On the other hand, the drought tolerant wild watermelon highly accumulates citrulline and CLMT2, a type 2 metallothionein, both with an extremely efficient hydroxyl radical scavenger activity, effectively protecting proteins and DNA from oxidative damage.<sup>37,80,81</sup>

**Photorespiration as a cellular protective route under drought.** Although photorespiration produces  $H_2O_2$  which is a potent thiol inhibitor, it is an extremely beneficial pathway for plants during

drought stress, when the rate of RuBP carboxylation is reduced due to the limitation of  $CO_2$  fixation by stomatal closure.<sup>15,35</sup> First of all it is an alternative route in energy dissipation which would otherwise provoke photo-inhibition of the photosynthetic apparatus.<sup>2</sup> Secondly it shifts  $H_2O_2$  production to peroxisomes rather than chloroplasts where the production of the hydroxyl radical is favored and where thiol enzymes of the Calvin cycle are targets to  $H_2O_2$  inhibition. Furthermore  $H_2O_2$  production in the peroxisomes is counterbalanced by the reductant-independent scavenging action of catalase as well as other peroxisomal APX isoforms which promptly detoxify the organelle. Finally, the photorespiration cycle yields glycine which is the precursor of glutathione, a metabolite of the ascorbate-glutathione pathway, hence providing additional protection against oxidative

stress.<sup>15,82</sup> Plant peroxisomes are dynamic organelles that adapt their number in response to environmental changes<sup>83</sup> and have been shown to proliferate in response to  $H_2O_2$  accumulation<sup>84</sup> which suggest that this is also likely to occur during drought stress.

**Mitochondria, alternative oxidase and ROS production avoidance.** The avoidance of ROS production during drought stress is also an important strategy that enables plants to cope with water shortage without extensive damage. Mitochondria play an important role in the avoidance of ROS production by efficient energy dissipation mechanisms.<sup>18</sup> The alternative oxidase (AOX) pathway is an alternative to the cytochrome pathway in the mitochondria and it diverts electrons flowing through the electron transport chain to produce water by the reduction of  $O_2$ .<sup>85,86</sup> In the case of durum wheat mitochondria, a Mediterranean plant well adapted to drought, three active energy-dissipating systems have been shown to coexist: the ATP-sensitive plant mitochondrial potassium channel (PmitoK<sub>ATP</sub>),<sup>87</sup> the plant uncoupling protein (PUCP)<sup>88</sup> and the alternative oxidase (AOX).<sup>89</sup> In fact, the activation of such energy-dissipating systems causes a significant reduction in mitochondrial ROS production.<sup>90,91</sup> Both the PmitoK<sub>ATP</sub> and the PUCP are modulated by the level of ROS production thus making them a very effective mechanism of ROS steady-state level regulation.<sup>91</sup> AOX protein level and capacity have also been shown to increase in the presence of  $H_2O_2$ .<sup>92,93</sup> On the other hand, AOX seems to be under the control of the photorespiratory cycle. In fact in durum wheat mitochondria it has been shown that the intermediate products of the photorespiratory pathway (glyoxylate and hydroxypyruvate) activate AOX,<sup>89,91</sup> hence establishing a cooperative mechanism of energy dissipation and ROS avoidance under drought stress when  $CO_2$  fixation is inhibited and photorespiration is at its maximum.

## ROS Signaling Under Drought

There is a frail balance between ROS production and scavenging that defines the normal steady-state level of intracellular ROS. Under drought stress this balance suffers an upward shift, ROS production being enhanced due to stomatal closure and the concomitant limitation on CO<sub>2</sub> fixation (Fig. 3). Instead of having an immediate deleterious effect, this raise in ROS production is likely to be beneficial to the plant if nevertheless kept under tight control. In fact, enhanced cellular ROS production is sensed by the plant as an alarm signal that triggers defense pathways and acclimatory responses, enabling the plant to adapt to the changing environment. Numerous reports highlight a signaling role for ROS production under stress.<sup>3,5,6,57,94-99</sup>

**H<sub>2</sub>O<sub>2</sub> as a secondary messenger.** The candidate ROS that is most likely to act as a secondary messenger in a stress-response signal transduction pathway is H<sub>2</sub>O<sub>2</sub>. The hydroxyl radical (HO•) has a diffusive-limited reaction due to its half-life (~1 μs) and its extremely reactive nature. In fact, it is impossible for this ROS to migrate from its site of production and function as a signal molecule, instead it will react locally with other molecules including itself and other ROS, proteins and lipids.<sup>100</sup> Regarding H<sub>2</sub>O<sub>2</sub>, it is not only the most stable ROS with the ability to easily diffuse from one cellular compartment to another but it also can be readily metabolized by an efficient cellular antioxidant system. Since it is produced at high rates under drought stress, a decrease in its concentration by the action of the antioxidant system allows for the rapid switch “on” and “off” of the signal, a condition essential for a secondary messenger to be effective. On the other hand, the relative toxicity of this active oxygen species has been put to question,<sup>3,14,16</sup> and its affinity to protein thiol groups suggests its possible role as a modulator of protein conformation and/or biochemical activities.<sup>101</sup> Furthermore, the intracellular thiol status is considered to be one of the mechanisms that enable for ROS sensing.<sup>101</sup> Plants are extremely tolerant of H<sub>2</sub>O<sub>2</sub> in comparison to animals and the antioxidant systems appear to function as tight controllers of the cellular redox-state instead of annihilators of all intracellular H<sub>2</sub>O<sub>2</sub>.<sup>14,101</sup> In this sense, antioxidants are key components on the modulation of the ROS signal since they determine the lifetime and the intensity of this signal.<sup>101,102</sup>

The specificity of the cellular ROS signal can be determined by its site of production, control and transduction.<sup>16</sup> Hence, the different plant cell compartments will influence differently the setting of the cellular redox signal under drought stress. Although the rate of H<sub>2</sub>O<sub>2</sub> production is faster in peroxisomes and chloroplasts,<sup>16</sup> mitochondria are the most vulnerable organelles to oxidative damage.<sup>103</sup> This can be explained by a lower antioxidant buffering in mitochondria as compared to peroxisomes and chloroplast. In this sense mitochondria play a crucial role in setting the cellular redox-state and initiating signal transduction cascades under drought stress.

Up to now, the known downstream events modulated by H<sub>2</sub>O<sub>2</sub> are calcium mobilisation, protein phosphorylation and gene expression.<sup>97</sup> Changes in cytosolic free calcium ([Ca<sup>2+</sup>]<sub>cyt</sub>) have been reported in numerous abiotic and biotic signal transduction pathways.<sup>104</sup> It has been shown that ROS induces an increase in [Ca<sup>2+</sup>]<sub>cyt</sub> by the activation of hyperpolarization-dependent Ca<sup>2+</sup>-permeable channels in the plasma membrane of *Arabidopsis* guard cells.<sup>105</sup> This ROS induced increase in [Ca<sup>2+</sup>]<sub>cyt</sub> has also been detected in other cell

types which suggests that the activation of Ca<sup>2+</sup> channels could be a key step in many ROS-mediated processes.<sup>100</sup> On the other hand, several reports have shown that H<sub>2</sub>O<sub>2</sub> induces mitogen-activated protein kinases (MAPKs), which are in turn implicated in several signal transduction cascades that modulate gene expression.<sup>106-109</sup>

H<sub>2</sub>O<sub>2</sub> stress response signaling pathways promote the accumulation of several cellular protectants that may act directly or indirectly in the regulation of the cellular redox-status, and consequently control the extent of the signal itself. For instance H<sub>2</sub>O<sub>2</sub> has been shown to induce the expression of the nuclear gene encoding for the mitochondrial alternative oxidase (AOX), Aox1.<sup>110</sup> In *Arabidopsis thaliana* cellular suspension culture four genes have been shown to be induced by H<sub>2</sub>O<sub>2</sub> by an RNA differential display approach.<sup>111</sup> Amongst the induced transcripts detected was a clone with sequence homology to a DNA damage repair protein (DRT112), one identical to serine/threonine protein kinase gene (APK2b), and one similar to an *Arabidopsis* late embryogenesis-abundant (LEA) protein homologue senescence-associated, SAG21 (the fourth gene presented no sequence homology to any known gene).<sup>111</sup> It is interesting to find that the detected H<sub>2</sub>O<sub>2</sub>-induced genes seem to be involved in cellular repair/protection mechanisms (DNA damage repair protein and LEA protein) or in the H<sub>2</sub>O<sub>2</sub> stress response signal transduction pathway (the serine/threonine protein kinase). Regarding the LEA proteins, these are characteristically expressed during the acquisition of desiccation tolerance in seeds but they have also been extensively associated to drought tolerance in many plant species.<sup>112</sup> Although their cellular functions are still not clearly understood, it has been shown that a citrus dehydrin CuCOR9 (LEA family D11) protects catalase activity under a freeze-thaw process,<sup>113</sup> prevents lipid peroxidation under cold stress<sup>114</sup> and was later shown to have radical scavenging properties,<sup>115</sup> making it a potent antioxidant that enables plants to cope with several abiotic stresses. A wider transcriptomics analysis of H<sub>2</sub>O<sub>2</sub> regulated genes in *Arabidopsis* by cDNA microarray technology was further undertaken and a total of 175 genes were identified as being H<sub>2</sub>O<sub>2</sub> responsive, 113 up-regulated and 62 down-regulated (the chip representing ~30% of the whole genome).<sup>116</sup> Amongst the up-regulated genes, 14 ESTs with known function were selected and assessed for expression studies by RNA blots. Wilting treatment by rapid desiccation induced the expression of several of the selected ESTs and the authors showed that this effect was partially mediated by H<sub>2</sub>O<sub>2</sub> in the case of the ESTs encoding calmoduline, a calcium-binding protein, the DREB2A transcription factor, the *Arabidopsis* MAP kinase ATMPK3, and a zinc finger protein.<sup>116</sup> This suggests that H<sub>2</sub>O<sub>2</sub> is likely to be a key component in the orchestration of plant drought stress responses, modulating Ca<sup>2+</sup> signaling, MAPK cascades and gene expression.

H<sub>2</sub>O<sub>2</sub> has also been recently shown to promote NO mediated activation of the proteasome complex in mammal endothelial cells which is involved in the degradation of oxidatively damaged proteins.<sup>117</sup> This H<sub>2</sub>O<sub>2</sub> triggered activation of proteases could be suggested to also occur in plants since it has been previously shown that drought stress induces a raise in cellular endoproteolytic activity.<sup>118</sup> Plants submitted to drought stress are also subjected to heat stress because of the reduced transpiration flux due to stomatal closure. Heat shock proteins which are molecular chaperones involved in the heat stress response have been shown to be induced by H<sub>2</sub>O<sub>2</sub>, which suggests its involvement in the heat stress signaling pathway.<sup>119</sup>

**ABA and ROS signaling under drought.** There seems to be an intricate relation between the hormone ABA and ROS. Drought stressed plants show enhanced accumulation of ABA and this triggers downstream responses that adapt the plant to the stress condition in an ABA-dependent manner.<sup>120-123</sup> However, several studies have shown that some ABA-dependent water stress responses cannot be elicited by ABA alone.<sup>99</sup> For instances, ABA accumulation is needed to induce proline accumulation in *Arabidopsis thaliana* at low water potentials ( $\Psi_w$ ) but exogenous application of ABA at high  $\Psi_w$  does not reproduce low  $\Psi_w$  induction of proline accumulation.<sup>124</sup> This suggests that other factors besides ABA are required to modulate the ABA response under water stress. One hypothesis that could explain this is the metabolic status of the plant.<sup>99</sup> In fact drought stress induces many physiological and biochemical changes that alter the metabolic status of the plant which could influence the cellular susceptibility to ABA accumulation. One important modification induced by drought stress is in the cellular redox-status by enhanced ROS production.<sup>101</sup> Hence ROS production under drought stress has been suggested to be the link between the metabolic status and ABA signaling that acts downstream of ABA and modulates the ABA signal transduction pathway.<sup>99</sup> ABA-dependent proline accumulation under drought stress highlights further the intricate and complex relation between ABA and ROS since the proposed functions of proline under stress are ROS scavenging<sup>125</sup> and regulation of the redox-status.<sup>126</sup>

One of the best characterized ABA-induced physiological responses under drought stress is leaf stomatal closure. This response operates more or less precociously on the onset of the drought period, depending on the plants' inherent strategy (and acclimation) towards drought stress. In the recent years major findings have shown that ABA activates the synthesis of  $H_2O_2$  in guard cells by a membrane bound NADPH oxidase and that  $H_2O_2$  mediates stomatal closure by activating (through hyperpolarization) plasma membrane  $Ca^{2+}$  channels.<sup>98,105,127</sup> Furthermore, the use of several *Arabidopsis* ABA mutants have enabled the dissection of some sequencing events in the pathways involving ABA and ROS signaling in guard cells. For instances, ABI1, and ABI2, two protein phosphatase 2C (PP2C)-like enzymes which are negative regulators of the ABA signal in guard cells, were suggested to act respectively up-stream and down-stream of ABA/ROS-mediated stomatal closure.<sup>127-129</sup> The sequencing events proposed in guard cells early ABA signal transduction were as follows: ABA, *abi1-1*, NAD(P)H-dependent ROS production, *abi2-1*, hyperpolarization-activated ( $I_{Ca}$ )  $Ca^{2+}$  channel activation followed by stomatal closing.<sup>127</sup> A different *Arabidopsis* mutant, *ost1* was like *abi1-1* impaired in ABA-induced ROS production and stomatal closure.<sup>130</sup> However, if external  $H_2O_2$  or calcium was applied, the mutant stomata responded like the wild-type. The protein OST1 is a Ser/Thr protein kinase which has a positive effect on ABA-mediated stomatal closure, acting upstream of ABA-induced ROS production.<sup>130</sup> Recent data have shown that *Arabidopsis thaliana* glutathione peroxidase 6 (ATGPX6) is also involved in ABA mediated guard cell  $H_2O_2$  signal transduction through a negative effect in ABI2 activity.<sup>131</sup> These authors also showed that transgenic plants overexpressing ATGPX6 were more resistant to drought stress and recovered better than the wild type. ATGPX6 was suggested to have a dual role, being not only an important scavenger of  $H_2O_2$ , but also an essential element of the ABA signaling pathway mediating stomatal regulation in response to drought stress.<sup>131</sup>

All this gathered places ROS in a central role in the guard cell ABA signaling network.<sup>132</sup> Furthermore, ROS signaling under drought stress acts not only downstream of stomatal closure but also upstream in the ABA signaling network. Interestingly, using wheat seedling root tips it has been shown that ROS (and NO) production plays a role in ABA synthesis under drought stress.<sup>133</sup>

**Sugars and ROS signaling.** Sugars, and more precisely soluble sugars such as glucose and sucrose, seem to play a dual role with respect to ROS, either promoting ROS production or participating indirectly in ROS scavenging mechanisms through NADPH generating pathways, such as the oxidative pentose-phosphate pathway.<sup>134</sup> Photosynthesis activity leads to the accumulation of ROS, by the Mehler reaction and also sugars. Furthermore, sugars have been shown to be involved in the regulation of the expression of several photosynthetic related genes as well as some ROS related genes such as superoxide dismutase.<sup>134,135</sup> Hence it has been suggested that soluble sugars could function as signals, useful for the plant in sensing and controlling not only the photosynthetic activity but also the cellular redox balance.<sup>134</sup> The connection between sugar sensing and ROS signaling is however extremely complex and seems to also involve ABA signaling, at least in some specific pathways. This can be illustrated by the ABA-induced proline accumulation under water deficit. Proline accumulation under water deficit is produced in an ABA-dependent manner. In a recent work, the use of several *Arabidopsis* ABA-insensitive mutants, (*abi1*, *abi2*, *abi3*, *abi4* and *abi5*) revealed that *abi4* had an increased proline accumulation under water deficit but presented a decreased sensitivity to exogenously applied ABA.<sup>124</sup> This response was altered by the supply of sucrose indicating that ABI4 has a role in connecting ABA and sugar in regulating proline accumulation.<sup>124</sup>

## Concluding Remarks and Future Challenges

For the last 15 years much information has been gathered regarding the involvement of a dynamic ROS equilibrium in the plant drought response. Although ROS can reach phytotoxic levels if drought stress is prolonged over to a certain extent (Fig. 3), the early signaling role of ROS under drought stress has been unequivocally established. The ROS signaling role is under tight modulation of the scavenging system and slight changes in ROS production or scavenging action under drought stress are likely to have immediate effects on the signal transduction. However, in order to have a clear view of the chronology of the events triggered by drought stress, there is a great need to homogenize drought treatments amongst researchers. In fact, it is quite difficult to compare the response of a plant submitted to a progressive drought stress, imposed by water withdrawal over several days, to the response of a plant submitted to a more immediate drought stress, imposed by watering with an osmotically active agent such as PEG for several hours. Furthermore, the response will also be quite different if the drought treatment is imposed on the intact plant, where the different organs cooperate as a whole or if it is applied to cut leaves. The terms "moderate" and "severe" drought stresses are also quite subjective and vary from one group to another. Recently, Verslues et al<sup>136</sup> have proposed several protocol-systems to study and quantify resistance to drought which could be extremely useful in a future standardization of water stress treatments.

Although the relationship between ROS, ABA, Ca<sup>2+</sup> and sugars has been revealed, it is not a straightforward one and the molecular nature of the interconnecting pathways remains to be solved. ABA, ROS and Ca<sup>2+</sup> are common players that are involved in cross-tolerance to many types of abiotic and biotic stress.<sup>95,137,138</sup> Nevertheless the information gathered shows only a glimpse on the intricate, complex and intriguing relation between ABA, Ca<sup>2+</sup> and ROS signaling that is still far from being completely resolved. Some fundamental questions remain unanswered: is ROS production enhancement under drought due mainly to “side effects” of stomatal closure or are there other oxidative bursts generated by positive feedback loops (like guard cell NADPH oxidases)? Are there cellular ROS receptors? How toxic really are ROS to plant cells? Does their toxicity change with the plants physiological status and/or acclimation? The use of transgenic plants overexpressing or expressing antisense constructs resulting in inhibition of specific scavenging enzymes, or the use of mutants with impaired H<sub>2</sub>O<sub>2</sub> generation will be extremely useful and are likely to help further in the study of the antioxidative mechanisms, and in disclosing the functions and biological roles of H<sub>2</sub>O<sub>2</sub> in response to drought stress.

#### Acknowledgements

I thank Christophe Bailly for the critical reading of an early version of the manuscript and José Santos for making the figures.

#### References

- Levitt J. Responses of plants to environmental stresses. New York: Academic Press, 1972.
- Smirnoff N. The role of active oxygen in the response of plants to water deficit and desiccation. *New Phytol* 1993; 125:27-58.
- Mittler R. Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci* 2002; 7:405-10.
- Levine A, Tenhaken R, Dixon R, Lamb C. H<sub>2</sub>O<sub>2</sub> from the oxidative burst orchestrates the plant hypersensitive disease resistance response. *Cell* 1994; 79:583-93.
- Dat J, Vandenberghe S, Vranová E, Van Montagu M, Inzé D, Van Breusegem F. Dual action of the active oxygen species during plant stress responses. *Cel Molec Life Sci* 2000; 57:779-95.
- Vranova E, Inzé D, Van Breusegem F. Signal transduction during oxidative stress. *J Exp Bot* 2002; 53:1227-36.
- Foyer CH, Harbinson J. Oxygen metabolism and the regulation of photosynthetic electron transport. In: Foyer CH, Mullineaux P, eds. Causes of Photooxidative Stresses and Amelioration of Defense Systems in Plants. Boca Raton, FL: CRC Press, 1994:1-42.
- Jakob B, Heber U. Photoproduction and detoxification of hydroxyl radicals in chloroplasts and leaves in relation to photoinactivation of photosystems I and II. *Plant Cell Physiol* 1996; 37:629-35.
- Noctor G, Foyer CH. Ascorbate and glutathione: Keeping active oxygen under control. *Annu Rev Plant Physiol Plant Mol Biol* 1998; 49:249-79.
- Foyer CH, Descourvières P, Kunert KJ. Protection against oxygen radicals: An important defence mechanism studied in transgenic plants. *Plant Cell Environ* 1994; 17:507-23.
- Baker NR. A possible role for photosystem II in environmental perturbations of photosynthesis. *Physiol Plant* 1991; 81:563-70.
- Asada K, Takahashi M. Production and scavenging of active oxygens in chloroplasts. In: Kyle DJ, Osmond CB, Arntzen CJ, eds. *Photoinhibition*. Amsterdam: Elsevier, 1987:227-87.
- Mehler AH. Studies on reactivity of illuminated chloroplasts. I. Mechanism of the reduction of oxygen and other Hill reagents. *Arch Biochem Biophys* 1951; 33:65-77.
- Foyer CH, Noctor G. Oxygen processing in photosynthesis: Regulation and signalling. *New Phytol* 2000; 146:359-88.
- Wingler A, Lea PJ, Quick WP, Leegood RC. Photorespiration: Metabolic pathways and their role in stress protection. *Phil Trans R Soc Lond B* 2000; 355:1517-29.
- Foyer CH, Noctor G. Redox sensing and signalling associated with reactive oxygen in chloroplasts, peroxisomes and mitochondria. *Physiol Plant* 2003; 119:355-64.
- Puntarulo S, Sanchez RA, Boveris A. Hydrogen peroxide metabolism in soybean embryonic axes at the onset of germination. *Plant Physiol* 1988; 86:626-30.
- Møller IM. Plant mitochondria and oxidative stress: Electron transport, NADPH turnover and metabolism of reactive oxygen species. *Annu Rev Plant Physiol Plant Mol Biol* 2001; 52:561-91.
- Bowler C, Van Montagu M, Inzé D. Superoxide dismutases and stress tolerance. *Annu Rev Plant Physiol Plant Mol Biol* 1992; 43:83-116.
- Willekens H, Chamnongpol S, Davey M, Schraudner M, Langebartels C, Van Montagu M, Inzé D, Van Camp W. Catalase is a sink for H<sub>2</sub>O<sub>2</sub> and is indispensable for stress defense in C3 plants. *EMBO J* 1997; 10:1723-32.
- Loewus FA. Ascorbic acid and its metabolic products. In: Preiss J, ed. *The Biochemistry of Plants*. Vol. 14. New York: Academic Press, 1988:85-107.
- Asada K. The water-water cycle in chloroplasts: Scavenging of active oxygens and dissipation of excess photons. *Annu Rev Plant Physiol* 1999; 50:601-39.
- Zhang J, Davies WJ. Increased synthesis of ABA in partially dehydrated root tips and ABA transport from roots to leaves. *J Exp Bot* 1987; 38:2015-23.
- Davies WJ, Zhang J. Root signals and the regulation of growth and development of plants in drying soil. *Annu Rev Plant Physiol Plant Mol Biol* 1991; 42:55-76.
- Hartung W, Sauter A, Hose E. Abscisic acid in xylem: Where does it come from, where does it go? *J Exp Bot* 2002; 53:27-32.
- Jiang F, Hartung W. Long-distance signalling of abscisic acid (ABA): The factors regulating the intensity of the ABA signal. *J Exp Bot* 2007; doi: 10.1093/jxb/erm127.
- Sgherri CLM, Pinzino C, Navari-Izzo F. Chemical changes and O<sub>2</sub> production in thylakoid membranes under water stress. *Physiol Plant* 1993; 87:211-6.
- Moran JF, Becana M, Iturbe-Ormaetxe I, Frechilla S, Klucas RV, Aparicio-Tejo P. Drought induces oxidative stress in pea plants. *Planta* 1994; 194:346-52.
- Sgherri CLM, Navari-Izzo F. Sunflower seedlings subjected to increasing water deficit stress: Oxidative stress and defense mechanisms. *Physiol Plant* 1995; 93:25-30.
- Wellburn FAM, Lau KK, Milling PMK, Wellburn AR. Drought and air pollution affect Nitrogen cycling and free radical scavenging in *Pinus halepensis* (Mill.). *J Exp Bot* 1996; 47:1361-7.
- Loggini B, Scartazza A, Brugnoli E, Navari-Izzo F. Antioxidative defense system, pigment composition, and photosynthetic efficiency in two wheat cultivars subjected to drought. *Plant Physiol* 1999; 119:1091-9.
- Boo YC, Jung J. Water deficit-induced oxidative stress and antioxidative defenses in rice plants. *J Plant Physiol* 1999; 155:255-61.
- Biehler K, Fock H. Evidence for the contribution of the Mehler-Peroxidase reaction in dissipating excess electrons in drought-stressed wheat. *Plant Physiol* 1996; 112:265-72.
- Sgherri CLM, Pinzino C, Navari-Izzo F. Sunflower seedlings subjected to increasing stress by water deficit: Changes in O<sub>2</sub> production related to the composition of thylakoid membranes. *Physiol Plant* 1996; 96:446-52.
- Noctor G, Veljovic-Jovanovic S, Driscoll S, Novitskaya L, Foyer C. Drought and oxidative load in the leaves of C<sub>3</sub> plants: A predominant role for photorespiration? *Ann Bot* 2002; 89:841-50.
- Asada K. Production and scavenging of reactive oxygen species in chloroplasts and their functions. *Plant Physiol* 2006; 141:391-6.
- Yokota A, Kawasaki S, Iwano M, Nakamura C, Miyake C, Akashi K. Citrulline and DRIP-1 protein (ArgE homologue) in drought tolerance of wild watermelon. *Ann Bot* 2002; 89:825-32.
- Pastori GM, Trippi VS. Oxidative stress induces high rate of glutathione reductase synthesis in a drought resistant maize strain. *Plant Cell Physiol* 1992; 33:957-61.
- Contour-Ansel D, Torres-Franklin ML, Cruz de Carvalho MH, D'Arcy-Lameta A, Zuily-Fodil Y. Glutathione reductase in leaves of cowpea: Cloning of two cDNAs, expression and enzymatic activity under progressive drought stress, desiccation and abscisic acid treatment. *Ann Bot* 2006; 98:1279-87.
- D'Arcy-Lameta, Ferrari-Iliou R, Contour-Ansel D, Pham-Thi AT, Zuily-Fodil Y. Isolation and characterization of four ascorbate peroxidase cDNAs responsive to water deficit in cowpea leaves. *Ann Bot* 2006; 97:133-40.
- Guo Z, Ou W, Lu S, Zhong Q. Differential responses of antioxidative system to chilling and drought in four rice cultivars differing in sensitivity. *Plant Physiol Biochem* 2006; 44:828-36.
- Lascano HR, Antonicelli GE, Luna CM, Melchiorre MN, Gomez LD, Racca RW, Trippi VS, Casano LM. Antioxidant system response of different wheat cultivars under drought: Field and in vitro studies. *Aust J Plant Physiol* 2001; 28:1095-102.
- Sairam PK, Saxena DC. Oxidative stress and antioxidants in wheat genotypes: Possible mechanism of water stress tolerance. *J Agron Crop Sci* 2000; 184:55-61.
- Edjolo A, Laffray D, Guerrier G. The ascorbate-glutathione cycle in the cytosolic and chloroplastic fractions of drought-tolerant and drought-sensitive poplars. *J Plant Physiol* 2001; 158:1511-7.
- Lima ALS, DaMatta FM, Pinheiro HA, Totola MR, Loureiro ME. Photochemical responses and oxidative stress in two clones of *Coffea canephora* under water deficit conditions. *Environ Exp Bot* 2002; 47:239-47.
- Turkan I, Bor M, Ozdemir F, Koca H. Differential responses of lipid peroxidation and antioxidants in the leaves of drought-tolerant *P. acutifolius* Gray and drought-sensitive *P. vulgaris* L. subjected to polyethylene glycol mediated water stress. *Plant Sci* 2005; 168:223-31.
- Torres-Franklin ML, Contour-Ansel D, Zuily-Fodil Y, Pham-Thi AT. Molecular cloning of glutathione reductase cDNAs and analysis of GR gene expression in cowpea and common bean leaves during recovery from a moderate drought stress. *J Plant Physiol* 2007; doi:10.1016/j.jplph.2007.03.011.
- Quartacci MF, Navari-Izzo F. Water stress and free radical mediated changes in sunflower seedlings. *J Plant Physiol* 1992; 139:612-25.
- Badiani M, De Biasi MG, Cologna M, Artemi F. Catalase, peroxidase and superoxide dismutase activities in seedlings submitted to increasing water deficit. *Agrochim* 1990; 34:90-102.
- Mittler R, Zilinskas BA. Regulation of pea cytosolic ascorbate peroxidase and other antioxidant enzymes during the progression of drought stress and following recovery from drought. *Plant J* 1994; 5:397-405.

51. Sharma P, Dubey RS. Drought induces oxidative stress and enhances the activities of anti-oxidant enzymes in growing rice seedlings. *Plant Grow Reg* 2005; 46:209-21.
52. Sofo A, Dichio B, Xiloyannis C, Masia A. Antioxidant defenses in olive trees during drought stress: Changes in activity of some antioxidant enzymes. *Funct Plant Biol* 2005; 32:45-53.
53. Cruz de Carvalho MH, Laffray D, Louguet P. Comparison of the physiological responses of *Phaseolus vulgaris* and *Vigna unguiculata* cultivars when submitted to drought conditions. *Environ Exp Bot* 1998; 40:197-207.
54. Perl A, Perl-Treves R, Galili S, Aviv D, Shalgi E, Malkin S, Galun E. Enhanced oxidative-stress defense in transgenic potato overexpressing tomato Cu,Zn superoxide dismutases. *Theor Appl Genet* 1993; 85:568-76.
55. Sen Gupta A, Heinen JL, Holaday AS, Burke JJ, Allen RD. Increased resistance to oxidative stress in transgenic plants that overexpress chloroplastic Cu / Zn superoxide dismutase. *Proc Natl Acad Sci USA* 1993; 90:1629-33.
56. Sen Gupta A, Webb RR, Holaday AS, Allen RD. Overexpression of superoxide dismutase protects plants from oxidative stress. *Plant Physiol* 1993; 103:1067-73.
57. Van Camp W, Capiu K, Van Montagu M, Inzé D, Slooten L. Enhancement of Oxidative stress tolerance in transgenic tobacco plants overproducing Fe-Superoxide dismutase in chloroplasts. *Plant Physiol* 1996; 112:1703-14.
58. Basu U, Good AG, Taylor GJ. Transgenic *Brassica napus* plants overexpressing aluminum-induced mitochondrial manganese superoxide dismutase cDNA are resistant to aluminum. *Plant Cell Environ* 2001; 24:1269-78.
59. Alscher RG, Ertuk N, Heath LS. Role of superoxide dismutases (SODs) in controlling oxidative stress in plants. *J Exp Bot* 2002; 372:1331-41.
60. McKersie BD, Bowley SR, Harjanto E, LepPrince O. Water-deficit tolerance and field performance of transgenic alfalfa overexpressing superoxide dismutase. *Plant Physiol* 1996; 111:1171-81.
61. Wang FZ, Wang QB, Kwon SY, Kwak SS, Su WA. Enhanced drought tolerance of transgenic rice plants expressing a pea manganese superoxide. *J Plant Physiol* 2005; 162:465-72.
62. Ratnayaka HH, Molin WT, Sterling TM. Physiological and antioxidant responses of cotton and spurred anoda under interference and mild drought. *J Exp Bot* 2003; 54:2293-305.
63. Keles Y, Oncel I. Response of the antioxidative defence system to temperature and water stress combinations in wheat seedlings. *Plant Sci* 2002; 163:783-90.
64. Dhindsa RS. Drought stress, enzymes of glutathione metabolism, oxidation injury, and protein synthesis in *Tortula ruralis*. *Plant Physiol* 1991; 95:648-51.
65. Rubio MC, González EM, Minchin FR, Webb KJ, Arrese-Igor C, Ramos J, Becana M. Effects of water stress on antioxidant enzymes of leaves and nodules of transgenic alfalfa overexpressing superoxide dismutases. *Physiol Planta* 2002; 115:531-40.
66. Jiang M, Zhang J. Water stress-induced abscisic acid accumulation triggers the increased generation of reactive oxygen species and upregulates the activities of antioxidant enzymes in maize leaves. *J Exp Bot* 2002; 53:2401-10.
67. Iturbe-Ormaetxe I, Escuredo Pedro R, Arrese-Igor C, Becana M. Oxidative damage in pea plants exposed to water deficit or paraquat. *Plant Physiol* 1998; 116:173-81.
68. Tang L, Kwon SY, Kim SH, Kim JS, Cho KY, Sung CK. Enhanced tolerance of transgenic potato plants expressing both superoxide dismutase and ascorbate peroxidase in chloroplasts against oxidative stress and high temperature. *Plant Cell Rep* 2006; 25:1380-6.
69. Lee SH, Ahsan N, Lee KW, Kim DH, Lee DG, Kwak SS, Kwon SY, Kim TH, Lee BH. Simultaneous overexpression of both CuZn superoxide dismutase and ascorbate peroxidase in transgenic tall fescue plants confers increased tolerance to a wide range of abiotic stresses. *J Plant Physiol* 2007; doi:10.1016/j.jplph.2007.01.003.
70. Anderson JV, Davis DG. Abiotic stress alters transcript profiles and activity of glutathione S-transferase, glutathione peroxidase, and glutathione reductase in *Euphorbia esula*. *Physiol Planta* 2004; 120:421-33.
71. Herbet P, Lenne C, Leblanc N, Julien JL, JoeDrevet R, Roedel-Drevet P. Two GPX-like proteins from *Lycopersicon esculentum* and *Helianthus annuus* are antioxidant enzymes with phospholipid hydroperoxide glutathione peroxidase and thioredoxin peroxidase activities. *Eur J Biol Chem* 2002; 269:2414-20.
72. Jung BG, Lee KO, Lee SS, Chi YH, Jang HH, Kang SS, Lee K, Lim D, Yoon SC, Yun D-J, Inoue Y, Cho MJ, Lee SY. A Chinese cabbage cDNA with high sequence identity to phospholipid hydroperoxide glutathione peroxidases encodes a novel isoform of thioredoxin-dependent peroxidase. *J Biol Chem* 2002; 277:12572-8.
73. Luna CM, Pastori GM, Driscoll S, Groten K, Bernard S, Foyer CH. Drought controls on H<sub>2</sub>O<sub>2</sub> accumulation, catalase (CAT) activity and CAT gene expression in wheat. *J Exp Bot* 2004; 56:417-23.
74. Zhang J, Kirkham MB. Antioxidant responses to drought in sunflower and sorghum seedlings. *New Phytol* 1992; 132:361-73.
75. Fu J, Huang B. Involvement of antioxidants and lipid peroxidation in the adaptation of two cool-season grasses to localized drought stress. *Environ Exp Bot* 2001; 45:105-14.
76. Shikanai T, Takedab T, Yamauchib H, Sanoa S, Tomizawaa KI, Yokotaa A, Shigeokab S. Inhibition of ascorbate peroxidase under oxidative stress in tobacco having bacterial catalase in chloroplasts. *FEBS Lett* 1998; 428:47-51.
77. Rizhsky L, Hallak-Herr E, Van Breusegem F, Rachmilevitch S, Barr JE, Rodermel S, Inzé D, Mittler R. Double antisense plants lacking ascorbate peroxidase and catalase are less sensitive to oxidative stress than single antisense plants lacking ascorbate peroxidase or catalase. *Plant J* 2002; 32:329-42.
78. Munné-Bosch S. The role of alpha-tocopherol in plant stress tolerance. *J Plant Physiol* 2005; 162:743-8.
79. Munné-Bosch S, Schwarz K, Alegre L. Enhanced formation of  $\alpha$ -tocopherol and highly oxidized abietane diterpenes in water stressed rosemary plants. *Plant Physiol* 1999; 121:1047-52.
80. Akashi K, Miyake C, Yokota A. Citrulline, a novel compatible solute in drought-tolerant wild watermelon leaves, is an efficient hydroxyl radical scavenger. *FEBS Lett* 2001; 508:438-42.
81. Akashi K, Nishimura N, Ishida Y, Yokota A. Potent hydroxyl radical-scavenging activity of drought-induced type-2 metallothionein in wild watermelon. *Biochem Biophys Res Comm* 2004; 323:72-8.
82. Noctor G, Arisi ACM, Jouanin L, Foyer CH. Photorespiratory glycine enhances glutathione accumulation in both the chloroplastic and the cytosolic compartments. *J Exp Bot* 1999; 50:1157-67.
83. Yan M, Rayapuram N, Subramani S. The control of peroxisome number and size during division and proliferation. *Curr Opin Cell Biol* 2005; 17:376-83.
84. Lopez-Huertas E, Charlton WL, Johnson B, Graham IA, Baker A. Stress induces peroxisome biogenesis genes. *EMBO J* 2000; 19:6770-7.
85. Vanlerberghe GC, McIntosh L. Alternative oxidase: From gene to function. *Annu Rev Plant Physiol Plant Mol Biol* 1997; 48:703-34.
86. Maxwell D, Wang Y, McIntosh L. The alternative oxidase lowers mitochondrial reactive oxygen. *Proc Natl Acad Sci USA* 1999; 96:8271-6.
87. Pastore D, Stoppelli MC, Di Fonzo N, Passarella S. The existence of the K<sup>+</sup> channel in plant mitochondria. *J Biol Chem* 1999; 274:26683-90.
88. Pastore D, Fratianni A, Di Pede S, Passarella S. Effect of fatty acids, nucleotides and reactive oxygen species on durum wheat mitochondria. *FEBS Lett* 2000; 470:88-92.
89. Pastore D, Trono D, Laus MN, Di Fonzo N, Passarella S. Alternative oxidase in durum wheat mitochondria: Activation by pyruvate, hydroxypyruvate and glyoxylate and physiological role. *Plant Cell Physiol* 2001; 42:1373-82.
90. Camacho A, Moreno-Sanchez R, Bernal-Lugo I. Control of superoxide production in mitochondria from maize mesocotyls. *FEBS Lett* 2004; 570:52-6.
91. Pastore D, Trono D, Laus MN, Di Fonzo N, Flagella Z. Possible plant mitochondria involvement in cell adaptation to drought stress. *J Exp Bot* 2007; 58:195-210.
92. Wagner AM. A role for active oxygen species as second messengers in the induction of alternative oxidase gene expression in *Petunia hybrida* cells. *FEBS Lett* 1995; 368:339-42.
93. Vanlerberghe GC, McIntosh L. Signals regulating the expression of the nuclear gene encoding alternative oxidase of plant mitochondria. *Plant Physiol* 1996; 111:589-95.
94. Foyer CH, Lopez-Delgado H, Dat J, Scott IM. Hydrogen peroxide- and glutathione-associated mechanisms of acclimatory stress tolerance and signalling. *Physiol Plant* 1997; 100:241-54.
95. Knight H, Knight MR. Abiotic stress signalling pathways: Specificity and cross-talk. *Trends Plant Sci* 2001; 6:262-7.
96. Neill SJ, Desikan R, Clarke A, Hurst RD, Hancock JT. Hydrogen peroxide and nitric oxide as signalling molecules in plants. *J Exp Bot* 2002; 53:1237-42.
97. Neill SJ, Desikan R, Hancock JT. Hydrogen peroxide signalling. *Curr Opin Plant Biol* 2002; 5:388-95.
98. Kwak JM, Mori IC, Pei Z-M, Leonhardt N, Torres MA, Dangel JL, Bloom RE, Bodde S, Jones JDG, Schroeder JL. NADPH oxidase AtrbohD and AtrbohF genes function in ROS dependent ABA signaling in Arabidopsis. *EMBO J* 2003; 22:2623-33.
99. Verslues PE, Zhu J-K. Before and beyond ABA: Upstream sensing and internal signals that determine ABA accumulation and response under abiotic stress. *Biochem Soc Trans* 2005; 33:375-9.
100. Mori IC, Schroeder JL. Reactive oxygen species activation of plant Ca<sup>2+</sup> channels. A signaling mechanism in polar growth, hormone transduction, stress signaling, and hypothetically mechanotransduction. *Plant Physiol* 2004; 135:702-8.
101. Foyer CH, Noctor G. Redox homeostasis and antioxidant signalling: A metabolic interface between stress perception and physiological responses. *Plant Cell* 2005; 17:1866-75.
102. Mittler R, Vanderauwera S, Gollery M, Van Breusegem F. Reactive oxygen gene network of plants. *Trends Plant Sci* 2004; 10:490-8.
103. Bartoli CG, Gomez F, Martinez DE, Guamet JJ. Mitochondria are the main target for oxidative damage in leaves of wheat (*Triticum aestivum* L.). *J Exp Bot* 2004; 403:1663-9.
104. Sanders D, Pelloux J, Brownlee C, Harper JF. Calcium at the crossroads of signaling. *Plant Cell* 2002; 14:S401-17.
105. Pei ZM, Murata Y, Benning G, Thomine S, Klusener B, Allen GJ, Grill E, Schroeder JL. Calcium channels activated by hydrogen peroxide mediate abscisic acid signalling in guard cells. *Nature* 2000; 406:731-4.
106. Desikan R, Clarke A, Hancock JT, Neill SJ. H<sub>2</sub>O<sub>2</sub> activates a MAP kinase-like enzyme in *Arabidopsis thaliana* suspension cultures. *J Exp Bot* 1999; 50:1863-6.
107. Grant JJ, Yun BW, Loake GJ. Oxidative burst and cognate redox signalling reported by luciferase imaging: Identification of a signal network that functions independently of ethylene, SA and Me-JA but is dependent on MAPKK activity. *Plant J* 2000; 24:569-82.
108. Kovtun Y, Chiu WL, Tena G, Sheen J. Functional analysis of oxidative stress-activated mitogen-activated protein kinase cascade in plants. *Proc Natl Acad Sci USA* 2000; 97:2940-5.
109. Samuel MA, Miles GP, Ellis BE. Ozone treatment rapidly activates MAP kinase signalling in plants. *Plant J* 2000; 22:367-76.
110. Maxwell D, Nickels R, McIntosh L. Evidence of mitochondrial involvement in the transduction of signals required for the induction of genes associated with pathogen attack and senescence. *Plant J* 2002; 29:269-79.

111. Desikan R, Neill SJ, Hancock JT. Hydrogen peroxide-induced gene expression in *Arabidopsis thaliana*. *Free Rad Biol Med* 2000; 28:773-8.
112. Ingram J, Bartels D. The molecular basis of dehydration tolerance in plants. *Annu Rev Plant Physiol Plant Mol Biol* 1996; 47:377-403.
113. Hara M, Terashima S, Kuboi T. Characterization and cryoprotective activity of cold-responsive dehydrin from *Citrus unshiu*. *J Plant Physiol* 2001; 158:1333-9.
114. Hara M, Terashima S, Fukaya T, Kuboi T. Enhancement of cold tolerance and inhibition of lipid peroxidation by citrus dehydrin in transgenic tobacco. *Planta* 2003; 217:290-8.
115. Hara M, Fujinaga M, Kuboi T. Radical scavenging activity and oxidative modification of citrus dehydrin. *Plant Physiol Biochem* 2004; 42:657-62.
116. Desikan R, A-H-Mackerness S, Hancock JT, Neill SJ. Regulation of the *Arabidopsis* transcriptome by oxidative stress. *Plant Physiol* 2001; 127:159-72.
117. Thomas S, Kotamraju S, Zielonka J, Harder DR, Kalyanaraman B. Hydrogen peroxide induces nitric oxide and proteasome activity in endothelial cells: A bell-shaped signaling response. *Free Rad Biol Med* 2007; 42:1049-61.
118. Cruz de Carvalho MH, d'Arcy-Lameta A, Roy-Macauley H, Gareil M, El-Maarouf H, Pham-Thi AT, Zuily-Fodil Y. Aspartic protease in leaves of common bean (*Phaseolus vulgaris* L.) and cowpea (*Vigna unguiculata* L. Walp): Enzymatic activity, gene expression and relation to drought susceptibility. *FEBS Lett* 2001; 492:242-6.
119. Volkov RA, Panchuk II, Mullineaux PM, Schöffl F. Heat stress-induced H<sub>2</sub>O<sub>2</sub> is required for effective expression of heat shock genes in *Arabidopsis*. *Plant Mol Biol* 2006; 61:733-746.
120. Shinozaki K, Yamaguchi-Shinozaki K. Gene expression and signal transduction in water-stress response. *Plant Physiol* 1997; 115:327-34.
121. Bray EA. Molecular responses to water deficit. *Plant Physiol* 1993; 103:1035-40.
122. Bray EA. Abscisic acid regulation of gene expression during water-deficit stress in the era of the *Arabidopsis* genome. *Plant Cell Environ* 2003; 25:153-61.
123. Zhu JK. Salt and drought stress signal transduction in plants. *Ann Rev Plant Biol* 2002; 53:247-73.
124. Verslues PE, Bray EA. Role of abscisic acid (ABA) and *Arabidopsis thaliana* ABA-insensitive loci in low water potential-induced ABA and proline accumulation. *J Exp Bot* 2006; 57:201-12.
125. Smirnoff N, Cumbes Q. Hydroxyl radical scavenging activity of compatible solutes. *Phytochem* 1989; 28:1957-60.
126. Hare PD, Cress WA, Van Staden J. Dissecting the roles of osmolyte accumulation during stress. *Plant Cell Environ* 1998; 21:535-53.
127. Murata Y, Pei ZM, Mori IC, Schroeder J. Abscisic acid activation of plasma membrane Ca<sup>2+</sup> channels in guard cells requires cytosolic NAD(P)H and is differentially disrupted upstream and downstream of reactive oxygen species production in *abi1-1* and *abi2-1* protein phosphatase 2C mutants. *Plant Cell* 2001; 13:2513-23.
128. Koornneef M, Reuling G, Karssen CM. The isolation and characterization of abscisic acid-insensitive mutants of *Arabidopsis thaliana*. *Physiol Plant* 1984; 61:377-83.
129. Gosti F, Bertauche N, Vartanian N, Giraudat J. Abscisic acid-dependent and -independent regulation of gene expression by progressive drought in *Arabidopsis thaliana*. *Mol Gen Genet* 1995; 246:10-8.
130. Mustilli AC, Merlot S, Vavasseur A, Fenzi F, Giraudat J. *Arabidopsis* OST1 protein kinase mediates the regulation of stomatal aperture by abscisic acid and acts upstream of reactive oxygen species production. *Plant Cell* 2002; 14:3089-99.
131. Miao Y, Lv D, Wang P, Wang XC, Chen J, Miao C, Song CP. An *Arabidopsis* glutathione peroxidase functions as both a redox transducer and a scavenger in abscisic acid and drought stress responses. *Plant Cell* 2006; 18:2749-66.
132. Desikan R, Cheung MK, Bright J, Henson D, Hancock JT, Neill SJ. ABA, hydrogen peroxide and nitric oxide signalling in stomatal guard cells. *J Exp Bot* 2004; 395:205-12.
133. Zhao Z, Chen G, Zhang C. Interaction between reactive oxygen species and nitric oxide in drought-induced abscisic acid synthesis in root tips of wheat seedlings. *Aust J Plant Physiol* 2001; 28:1055-61.
134. Coué I, Sulmon C, Gouesbet G, El Amrani A. Involvement of soluble sugars in reactive oxygen species balance and responses to oxidative stress in plants. *J Exp Bot* 2006; 57:449-59.
135. Sulmon C, Gouesbet G, El Amrani A, Coué I. Sugar induced tolerance to the herbicide atrazine in *Arabidopsis* seedlings involves activation of oxidative and xenobiotic stress responses. *Plant Cell Rep* 2006; 25:489-98.
136. Verslues PE, Agarwal M, Katiyar-Agarwal S, Zhu J, Zhu JK. Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. *Plant J* 2006; 45:523-39.
137. Bowler C, Fluhr R. The role of calcium and activated oxygens as signals for controlling cross-tolerance. *Trends Plant Sci* 2000; 5:241-5.
138. Fujita M, Fujita Y, Noutoshi Y, Takahashi F, Narusaka Y, Yamaguchi-Shinozaki K, Shinozaki K. Crosstalk between abiotic and biotic stress responses: A current view from the points of convergence in the stress signaling networks. *Curr Opin Plant Biol* 2006; 9:436-42.
139. Tian XR, Lei YB. Physiological responses of wheat seedlings to drought and UV-B radiation. Effect of exogenous sodium nitroprusside application. *Russ J Plant Physiol* 2007; DOI 10.1134/S1021443707050160.
140. Lei Y, Yin C, Li C. Differences in some morphological, physiological, and biochemical responses to drought stress in two contrasting populations of *Populus przewalskii*. *Physiol Planta* 2006; 127:182-91.
141. Pinheiro HA, DaMatta FM, Chaves ARM, Fontes EPB, Loureiro ME. Drought tolerance in relation to protection against oxidative stress in clones of *Coffea canephora* subjected to long-term drought. *Plant Sci* 2004; 167:1307-14.
142. Chen KM, Gong HJ, Chen GC, Wang SM, Zhang CL. Gradual drought under field conditions influences the glutathione metabolism, redox balance and energy supply in spring wheat. *J Plant Grow Reg* 2004; 23:20-8.